

# INORT: Biofertilizer based on *Inula viscosa* L. (*Dittrichia viscosa* L.), algae and microorganisms for growth, *Fusarium oxysporum* defence and water stress resistance of *Plumeria frangipani*

Domenico Prisa <sup>1,\*</sup> and Francesco Attanasio <sup>2</sup>

<sup>1</sup> CREA Research Centre for Vegetable and Ornamental Crops, Council for Agricultural Research and Economics, Via dei Fiori 8, 51012 Pescia, PT, Italy.

<sup>2</sup> Attanasio Farm, Via Matteotti 35, San Sebastiano al Vesuvio, NA, Italy.

International Journal of Biological and Pharmaceutical Sciences Archive, 2023, 06(02), 047–053

Publication history: Received on 28 August 2023; revised on 06 October 2023; accepted on 09 October 2023

Article DOI: <https://doi.org/10.53771/ijbpsa.2023.6.2.0097>

---

## Abstract

**Research objective:** In this study, the possibility of using INORT, a biostimulant based on *Inula viscosa* L. (*Dittrichia viscosa* L.), algae and microorganisms to improve growth, defence against *Fusarium oxysporum* and resistance to water stress, on *Plumeria frangipani* plants grown in open field was evaluated.

**Materials and Methods:** The experiments, started in January 2023, were conducted in the greenhouses of CREA-OF in Pescia (Pt), on *Plumeria frangipani* plants cvs “California sunset and “Super round”. The plants were grown in open field. The five experimental groups in cultivation were: group without biostimulant (CTRL); group with microorganisms; group with algae; group with *Inula viscosa*; group with (INORT) (mix Microorganisms + *Inula viscosa* + Algae). The product INORT was supplied by the Francesco Attanasio farm. On 15 September 2023, plant height, number of leaves, number of branches, total leaf area per plant (mm<sup>2</sup>), primary root length (mm), aerial and root system biomass were recorded. In addition, plant mortality as a result of *Fusarium oxysporum* attacks and water stress was assessed in the experiment.

**Results and Discussion:** The experiment showed that the use of the biostimulant based on *Inula viscosa*, microorganisms and algae (INORT) can indeed significantly improve the vegetative and root growth of *Plumeria frangipani* plants grown in the open field. Improvements were also found in plant height, number of leaves, number of floral branches, leaf area, vegetative and root biomass and root length at different irrigation frequencies of 3 and 6 days. A very interesting aspect was also the ability of the biostimulant based on *Inula viscosa*, microorganisms and algae to significantly reduce the incidence of *Fusarium oxysporum* attacks on leaves.

**Conclusions:** Given the importance of *Inula viscosa* from a medicinal, pollination and biodiversity point of view, new agricultural experiments are very important as they could allow the development of new biofertiliser products that can be used in organic and sustainable farming systems

**Keywords:** Microorganisms; Sustainable applications; Plant extract; Rhizosphere; Plant stimulation

---

## 1. Introduction

A perennial bushy plant (*Inula viscosa* L., also known as *Dittrichia viscosa* Greuter) belonging to the family *Compositae*, it is considered the largest among Phanerogams and includes about 959 genera, which are commonly found in Mediterranean areas [1]. An erect cauli, woody at the base, richly covered with leaves, and a height of 50 to 150 cm, *I. viscosa* has a characteristic smell, with erect cauli, woody at their base and thickly covered in leaves. Inflorescences

---

\* Corresponding author: Domenico Prisa

consist of numerous pyramidal flower heads with golden yellow flowers with linear-lanceolate leaves. In autumn, flowers bloom, and fruits are made up of achenes. Many studies have been conducted in relation to the presence of numerous compounds of different chemical nature in different parts of the plant (roots, stems, leaves, flowers, etc.), whose properties make some potential applications in pharmaceuticals, cosmetics, aromatics, and related fields [2]. Folk medicine uses this species primarily to treat liver disorders, including analgesic, anti-inflammatory, antipyretic, anthelmintic, and antifungal properties [3,4]. Although it is described as a typical plant of the Mediterranean region [5,6,7], it is actually found in southern Europe and North Africa, but no precise habitats of growth have been identified. It is very easy to recognize in rural environments and near roadsides, prefers calcareous and clayey soil, often acidic and being a heliophilous species, it can be found at a range of elevations from 0 to 800 m.a.s.l. For several years no scientific studies were conducted on its potential, but recently it has been revalued [8]. As a plant abundantly foraged by bees, *I. viscosa* provides a source of nourishment for butterfly caterpillars and moths [9,10]. Furthermore, it is recognized for its high melliferous power. *Inulas* bloom very long between late spring and early autumn, when most plants are generally no longer in bloom. As a result, the classroom has a high potential for producing swarms in the following spring, as well as honey in the autumn and late summer [11,12].

## 2. Related Work

Several studies have reported that *I. viscosa* extracts have the ability to control pathogenic fungi, while secondary plant metabolites have been shown to be insecticidal and antibacterial [14,16]. As a result of several studies it has been demonstrated that *Inula viscosa* plants play a role agro-ecological very important because: i) show high allelopathic activity as they produce chemical compounds toxic to other plant species; ii) act as a dwelling place for antagonists useful to combat plant pests and therefore fundamental in the biological struggle; iii) also find application in the phyto-remediation of contaminated soil in processes of phyto-extraction [17,18], bioaccumulation [19] and as bioindicator [20]. These activities are closely related to the phenological phases of the plant. Experimental evidence shows that *Inula* extract can be used in the control of varroa and nosema in bees, introduced into gel blocks and exploiting olfactory disorientation [21,22,23].

In this study, the possibility of using INORT, a biostimulant based on *Inula viscosa* L. (*Dittrichia viscosa* L.), algae and microorganisms to improve growth, defence against *Fusarium oxysporum* and resistance to water stress, on *Plumeria frangipani* plants grown in open field was evaluated.



**Figure 1** Details of the *Plumeria frangipani* plants used in the trial

## 3. Materials and methods

The experiments, started in January 2023, were conducted in the greenhouses of CREA-OF in Pescia (PT), on *Plumeria frangipani* plants cvs “California sunset” and “Super round” (Figure 1).

The plants were grown in open field; 30 rooted cuttings per thesis, divided into 3 replications of 10 plants each planted in early January 2023. All plants were fertilised with a slow-release fertiliser (2 kg m<sup>-3</sup> of Osmocote Pro® for 6 months) introduced into the growing medium at the time of transplanting.

. The five experimental groups in cultivation were:

- Group without biostimulant (CTRL): (peat 70% + pumice 20%), irrigated with water and substrate previously fertilized;
- Group with microorganisms (MC): (peat 70% + pumice 20%), irrigated with water and substrate previously fertilized, (Lactic acid bacteria, Photosynthetic bacteria, Yeast, Arbuscular mycorrhizae:  $4 \times 10^3$  spores/ml) 1% per week during the growing cycle (25 ml per plant);
- Group with algae (AG): (peat 70% + pumice 20%), irrigated with water and substrate previously fertilized, 1% per week during the growing cycle (25 ml per plant);
- Group with *Inula viscosa* (INU): (peat 70% + pumice 20%), irrigated with water and substrate previously fertilized, 1% per week during the growing cycle (25 ml per plant);
- Group with (INORT) (mix Microorganisms + *Inula viscosa* + Algae): (peat 70% + pumice 20%), irrigated with water and substrate previously fertilized, 1% per week during the growing cycle (25 ml per plant). The product INORT was supplied by the Francesco Attanasio farm.

The plants were watered once every 3 days and 6 days (two treatments with different irrigation intervals for each thesis to assess any difference in water stress resistance) and cultivated for 9 months. The plants were drip-irrigated. Irrigation was activated by a timer whose schedule was adjusted weekly according to the weather conditions and the leaching fraction. On 15 September 2023, plant height, number of leaves, number of branches, total leaf area per plant ( $\text{mm}^2$ ), primary root length (mm), aerial and root system biomass were recorded. In addition, plant mortality as a result of *Fusarium oxysporum* attacks and water stress was assessed in the experiment.

### 3.1. Statistics

The experiment was carried out in a randomized complete block design. Collected data were analysed by one-way ANOVA, using GLM univariate procedure, to assess significant ( $P \leq 0.05$ , 0.01 and 0.001) differences among treatments. Mean values were then separated by LSD multiple-range test ( $P = 0.05$ ). Statistics and graphics were supported by the programs Costat (version 6.451) and Excel (Office 2010).

## 4. Results and Discussion

The experiment showed that the use of the biostimulant based on *Inula viscosa*, microorganisms and algae (INORT) can indeed significantly improve the vegetative and root growth of *Plumeria frangipani* plants grown in the open field (Table 1 and Table 2). All treatments showed a significant improvement over the untreated control for the agronomic parameters analysed, but the INORT treatment was statistically the best (Figure 2). Improvements were also found in plant height, number of leaves, number of floral branches, leaf area, vegetative and root biomass and root length at different irrigation frequencies of 3 and 6 days (Figure 3). A very interesting aspect was also the ability of the biostimulant based on *Inula viscosa*, microorganisms and algae to significantly reduce the incidence of *Fusarium oxysporum* attacks on leaves.

Typically heliophilous and ruderal, *Inula viscosa* is commonly found in uncultivated areas, ruins, along roads, headlands, cliffs and escarpments. Its ruggedness and adaptability allow it to colonise poor, dry and stony soils [24,25]. It has sticky leaves and an odour that repels livestock. When invasive, it tends to grow in degraded pastures and in the rows of extensive plantations (vineyards, olive groves, orchards), but avoids arable land that is regularly cultivated. The plant was used in some areas of Sardinia as a remedy for rheumatic pains [26]. In Sicily it was used for healing and haemostatic purposes. In Tuscany, fresh leaves were used against excessive sweating. In Liguria, dried leaves were used by the poor as a substitute for tobacco. In barns it was mixed with hay to repel mice. As a melliferous plant, it is highly favoured by bees, mainly because of its abundance of pollen and its long flowering period. As a result, this plant produces multifloral honey in late summer and autumn, and monofloral honey in areas of high abundance. Inulas are attacked by a tephritid gall dipteran called *Myopites stylatus* [27,28]. The olive fly is the overwintering host of *Eupelmus urozonus*, a polyphagous parasitoid of *Hymenoptera calcidoides* that reproduces 2-3 times per year [29,30]. Integrated pest management programmes can contribute to the control of the phytophage by spreading *Eupelmus*, inula in uncultivated areas of olive groves, which is essential as *Eupelmus* is the most active natural antagonist of the olive fly [31,32]. In particular, *Polistes gallicus*, a species with a typical Mediterranean distribution, nests on the most developed branches of the plant [33,34]. *Aphis nerii* and *Varroa* appear to be susceptible to the toxic effects of costic acid. The antifungal activity of *I. viscosa* extracts against phytopathogenic fungi and secondary metabolites has been demonstrated in numerous studies, as well as its insecticidal, antifungal, acaricidal, antibacterial and cytotoxic properties [35]. In addition, biofertilising activity has been demonstrated in combination with microorganisms and algae on a variety of vegetable species [36]. Biostimulative effects have also been demonstrated in field-grown plants, where plants are more

resistant to water stress and to pathogenic fungi such as *Fusarium oxysporum*. *Inula viscosa* has a number of interesting properties that make it a potential liquid biostimulant and an effective means of controlling pathogenic insects and fungi.

**Table 1** Evaluation of *Inula viscosa*, algae and microorganisms on agronomic characters on plants of *Plumeria frangipani* cv California Sunset

California Sunset	Rounds (days)	PH (cm)	LN (n°)	NB (n°)	TLA (mm <sup>2</sup> )	VW (g)	RW (g)	RL (cm)	FO attack (n°)
CTRL	3	52.57 d	31.22 c	1.41 c	122.24 d	76.46 d	51.94 d	3.65 e	5.61 a
MC	3	59.38 b	34.23 b	2.22 b	124.99 c	78.18 c	53.84 c	4.13 d	3.63 b
AG	3	59.39 b	34.81 b	2.42 b	126.10 b	79.02 b	54.61 b	5.09 b	3.82 b
INU	3	55.82 c	34.82 b	2.43 b	124.75 c	77.96 c	53.73 c	4.81 c	2.84 c
INORT	3	61.74 a	38.00 a	3.61 a	132.17 a	80.52 a	56.61 a	5.47 a	1.00 d
ANOVA	-	***	***	***	***	***	***	***	***
CTRL	6	42.76 d	21.81 d	1.20 b	75.10 d	48.36 e	35.31 d	3.12 e	6.82 a
MC	6	48.28 b	25.00 b	2.00 a	79.40 b	50.48 c	38.57 b	4.35 b	3.83 b
AG	6	48.44 b	24.80 b	2.22 a	79.51 b	51.40 b	38.68 b	4.17 c	3.64 b
INU	6	45.77 c	23.00 c	1.41 b	77.84 c	49.37 d	37.57 c	3.76 d	4.22 b
INORT	6	49.55 a	27.61 a	2.22 a	83.93 a	52.27 a	49.39 a	5.02 a	2.21 c
ANOVA	-	***	***	**	***	***	***	***	***


One-way ANOVA; n.s. – non significant; \*, \*\*, \*\*\* – significant at  $P \leq 0.05$ , 0.01 and 0.001, respectively; different letters for the same element indicate significant differences according to Tukey's (HSD) multiple-range test ( $P = 0.05$ ). Parameters: PH = plant height (cm); LN = leaves number (cm); NB = branches number (n°); TLA = total leaves area (mm<sup>2</sup>); VW = vegetative weight (g); RW = roots weight (g); RL = roots length (cm); FO= plants affected by *Fusarium oxysporum* (n°). Treatments: CTRL=control; MC=microorganisms; INU=*Inula viscosa*; INORT=*Inula viscosa*+microorganisms+algae; AG= algae

**Table 2** Evaluation of *Inula viscosa*, algae and microorganisms on agronomic characters on plants of *Plumeria frangipani* cv Super Round


Super Round	Rounds (days)	PH (cm)	LN (n°)	NB (n°)	TLA (mm <sup>2</sup> )	VW (g)	RW (g)	RL (cm)	FO attack (n°)
CTRL	3	52.59 d	32.41 c	1.82 c	121.89 d	76.48 d	51.97 d	3.69 e	6.00 a
MC	3	59.39 b	34.82 b	2.42 bc	125.21 c	78.27 c	53.86 c	4.28 d	3.82 b
AG	3	59.41 b	35.00 b	2.61 b	126.14 b	79.06 b	56.64 b	5.12 b	3.43 b
INU	3	55.82 c	35.00 b	2.61 b	124.76 c	78.16 c	53.74 c	4.82 c	2.61 c
INORT	3	61.75 a	38.21 a	3.82 a	132.77 a	80.63 a	56.84 a	5.48 a	0.64 d
ANOVA	-	***	***	***	***	***	***	***	***
CTRL	6	42.77 d	22.22 d	1.42 c	75.31 d	48.39 e	35.32 d	3.17 d	7.42 a
MC	6	48.29 b	25.42 b	2.21 ab	79.58 b	50.56 c	38.66 b	4.37 b	3.61 b
AG	6	48.45 b	25.00 b	2.63 a	79.56 b	51.51 b	38.73 b	4.26 b	3.23 b
INU	6	45.78 c	23.41 c	1.63 bc	77.85 c	49.46 d	37.59 c	3.77 c	3.42 b
INORT	6	49.56 a	28.41 a	2.82 a	84.08 a	52.43 a	43.66 a	5.11 a	1.64 c

ANOVA	-	***	***	***	***	***	***	***	***
-------	---	-----	-----	-----	-----	-----	-----	-----	-----

One-way ANOVA; n.s. – non significant; \*, \*\*, \*\*\* – significant at  $P \leq 0.05$ , 0.01 and 0.001, respectively; different letters for the same element indicate significant differences according to Tukey's (HSD) multiple-range test ( $P = 0.05$ ); Parameters: PH = plant height (cm); LN = leaves number (cm); NB = branches number (n°); TLA = total leaves area (mm<sup>2</sup>); VW = vegetative weight (g); RW = roots weight (g); RL = roots length (cm); FO= plants affected by *Fusarium oxysporum* (n°). Treatments: CTRL=control; MC=microorganisms; INU=Inula viscosa; INORT=Inula viscosa+microorganisms+algae; AG= algae



**Figure 2** Effect of INORT on vegetative biomass, plant height, number of branches of *Plumeria frangipani* cv “California Sunset”. Turnover 3-day water regime. Legend: (CTRL): control; (INORT):Inula+microorganisms+algae



**Figure 3** Effect on leaf resistance to water stress of treatment with INORT on *Plumeria frangipani* cv Super round”. Irrigation turn-over of 6 days. Legend: (CTRL) control; (MC): microorganisms; (AG): algae; (INORT): microorganisms+algae+inula

## 5. Conclusion

The trial showed that the use of a liquid biostimulant based on *Inula viscosa* was able to significantly improve the growth, vegetative and root biomass of *Plumeria frangipani*. The treatment also provides increased resistance to water stress and protection against plant attack by *Fusarium oxysporum*.

This highlights other interesting and innovative aspects of the use of this plant, which have already been highlighted in previous trials on various potted plant species. Given the importance of *inula viscosa* from a medicinal, pollination and

biodiversity point of view, new agricultural experiments are very important as they could allow the development of new biofertiliser products that can be used in organic and sustainable farming systems.

---

## Compliance with ethical standards

### *Acknowledgments*

The research is part of the project "INORT": evaluation of derivatives of *Inula viscosa* in the growth and defense of horticultural plants. I would like to thank the Attanasio company for their cooperation, particularly in the person of Dr. Francesco Attanasio, who actively collaborated on the INORT project.

### *Disclosure of conflict of interest*

The author declares no conflict of interest.

### *Statement of ethical approval*

The present research work does not contain any studies performed on animal/humans subjects.

---

## References

- [1] Searchinger T. The Great Balancing Act: Installment 1 of Creating a Sustainable Food Future, WRI, Washington, D.C, 2013.
- [2] Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, Mueller ND, O'Connell C, Ray DK, West PC, Balzer C, Bennett EM, Carpenter SR, Hill J, Monfreda C, Polasky S, Rockström J, Sheehan J, Siebert S, Tilman GD, Zaks DPM. Solutions for a cultivated planet, *Nature*. 2011; 478: 337.
- [3] Parajuli R, Thoma G, Matlock MD. Environmental sustainability of fruit and vegetable production supply chains in the face of climate change: a review, *Sci. Total Environ*. 2019; 650: 2863–2879.
- [4] Zulfiqar F, Allaire SE, Akram NA, Méndez A, Younis A, Peerzada AM, Shaukat N, Wright S.R. Challenges in organic component selection and biochar as an opportunity in potting substrates: a review, *J. Plant Nutr*. 2019; 42: 1386–1401.
- [5] Prisa D. Improving Quality and Production of Horticultural Crops Through the Use of A Biostimulant Based on *Inula viscosa* and Control of Seedling Pathogens. *International Journal of Scientific Research in Biological Sciences*. 2021; 8(1): 21–27
- [6] Prisa D. Biofertilizer based on liquid fermented with *Inula viscosa*, microorganisms and algae in the growth and biocontrol of *Sphaerotheca pannosa* var. *rosae* of seed rose plants. *World Journal of Biology Pharmacy and Health Sciences*. 2021; 06(03): 020–026.
- [7] Prisa D, Gobbino M. (2021). Sustainable Methods Based on Microbial Biofertilizers and Plant Repellent Extracts in the Cultivation of Aloe Vera. *World Journal of Environmental Biosciences*. 2021; 10(4): 27–32
- [8] Mullins M.G. *Plant Improvement in Horticulture: The Case for Fruit Breeding*, The Regional Institute. 2006; 85–92.
- [9] Posmyk MM, Szafrańska K. Biostimulators: a new trend towards solving an old problem, *Front. Plant Sci*. 2016; 7: 7–48.
- [10] Kauffman GL, Kneivel DP, Watschke TL. Effects of a biostimulant on the heat tolerance associated with photosynthetic capacity, membrane thermostability, and polyphenol production of perennial ryegrass, *Crop Sci*. 2007; 47: 261–267.
- [11] Rady MM, Desoky ES, Elrys AS, Boghdady MS. Can licorice root extract be used as an effective natural biostimulant for salt-stressed common bean plants? *S. Afr. J. Bot*. 2019; 121: 294–305.
- [12] Prisa D. Germination Of Vegetable And Grassland species With Micronized chabazitic-Zeolites And Endophytic Fungi. *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)*. 2019; 12(51): 32–37.
- [13] Prisa D. Improvement Quality and Content of Pepper and Chilli Nitrates Influenced by the Effective Microorganisms. *American Scientific Research Journal for Engineering, Technology, and Sciences (ASRJETS)*. 2019; 53(1):176–181
- [14] Attanasio F, Prisa D. Plant extracts from mediterranean vegetation in the cultivation and defence of *Plectranthus coleoides* and *Plectranthus amboinicus*. *World Journal of Advanced Research and Reviews*, 2023; 18(02): 502–509.

- [15] Hussein NM, Hussein MI, Gadel Hak SH, Hammad MA. Effect of two plant extracts and four aromatic oils on tuta absoluta population and productivity of tomato cultivar gold stone, *Nat. Sci.* 2014; 12: 108–118.
- [16] Hayat S, Ahmad H, Ali M, Hayat K, Khan M, Cheng Z. Aqueous garlic extract as a plant biostimulant enhances physiology, improves crop quality and metabolite abundance, and primes the defense responses of receiver plants, *Appl. Sci.* 2018; 8: 1505.
- [17] Younis A, Akhtar MS, Riaz A, Zulfiqar F, Qasim M, Farooq A, Tariq U, Ahsan M, Bhatti Z.M. Improved cut flower and corm production by exogenous moringa leaf extract application on gladiolus cultivars, *Acta Sci. Pol.* 2018; 17: 25–38.
- [18] Prisa D, Attanasio F. Biostimulant derived from the fermentation of *Inula viscosa* (Inort) in the germination and growth of *Amaranthus hypochondriacus*. *World Journal of Advanced Research and Reviews.* 2022; 16(01): 027–033
- [19] Prisa D. Increasing plant resistance with silicon applications. *World Journal of Advanced Research and Reviews.* 2023; 17(03): 602–608.
- [20] Culver M, Fanuel T, Chiteka AZ. Effect of moringa extract on growth and yield of tomato, *Green. J. Agric. Sci.* 2012; 2: 207–211.
- [21] Nasir M, Khan AS, Basra SA, Malik AU. Foliar application of moringa leaf extract, potassium and zinc influence yield and fruit quality of ‘Kinnow’ mandarin, *Sci. Hortic.* 2016; 210: 227–235.
- [22] Aslam M, Sultana B, Anwar F, Munir H. Foliar spray of selected plant growth regulators affected the biochemical and antioXidant attributes of spinach in a field experiment, *Turk. J. Agric. For.* 2016; 40: 136–145.
- [23] Hemalatha M, Selvi D, Vasanthi D. Integrated plant nutrients supply and foliar nutrition on fruit yield, nutrient uptake and availability in okra [*Abelmoschus esculentus* (L.) Moench] hybrid COBH-4 in an alfisol of tamirabarani command area, *Int. J. Adv. Agric. Sci. Technol.* 2018; 5: 47–66.
- [24] Sakr WRA, El-Sayed AA, Hammouda AM, Saad El Deen FSA. Effect of NPK, aloe gel and moringa extracts on geranium plants, *J. Hortic. Sci. Ornam. Plants.* 2018; 10: 01–16.
- [25] Prisa D. *Ascophyllum nodosum* extract on growth plants in *Rebutia heliosa* and *Sulcorebutia canigueralli*. *GSC Biological and Pharmaceutical Sciences.* 2020; 10(01): 039–045
- [26] Abdou MAH, Badran FS, Ahmed ET, Taha RA, Abdel-Mola MAM. Effect of compost and some natural stimulant treatments on: ii. corms production and chemical constituents of (*Gladiolus grandiflorusc*v. Peter pears) plants, *Sci. J. Flowers Ornam. Plants.* 2018; 5: 115–126.
- [27] Ali M, Zhi-hui C, Hayat S, Ahmad H, Ghani MI, Tao L. Foliar spraying of aqueous garlic bulb extract stimulates growth and antioXidant enzyme activity in eggplant (*Solanum melongena* L.), *J. Integr Agric.* 2018; 17: 60345–60347.
- [28] Hanafy MS, Saadawy FM, Milad SMN, Ali R.M. Effect of some natural extracts on growth and chemical constituents of *Schefflera arboricola* plants, *J. Hortic. Sci. Ornam. Plants.* 2012; 4: 26–33.
- [29] Shakir MA, Al-Rawi WAA. Effect of garlic and licorice root extract on leaves mineral and hormonal content of pear transplants, *Iraqi J. Agric. Sci.* 2017; 48: 138.
- [30] Hayat S, Ahmad H, Ali M, Ren K, Cheng Z. Aqueous garlic extract stimulates growth and antioXidant enzymes activity of tomato (*Solanum lycopersicum*), *Sci. Hortic.* 2018; 240: 139–146.
- [31] Hafez OM, Saleh MA, El-Lethy SR. Response of some seedlings olive cultivars to foliar spray of yeast and garlic extracts with or without vascular arbuscular mycorrhizal fungi, *J. World Appl. Sci.* 2013; 24: 1119–1129.
- [32] El-Azim A, Rania MR, Badawy MYM. Effect of bio-fertilization and different licorice extracts on growth and productivity of *Foeniculum vulgare*, *Mill. Plant Middle East J. Agric.* 2017; 6: 1–12.
- [33] Sánchez-Gómez R, Garde-Cerdán T, Zalacaín A, Garcia R, Cabrita MJ, Salinas MR. Vine-shoot waste aqueous extract applied as foliar fertilizer to grapevines: effect on amino acids and fermentative volatile content, *Food Chem.* 2016; 197: 132–140.
- [34] Abdou MA, El-Sayed AA, Taha RAI, Sayed MAAE, Botros WSE. Effect of compost and some biostimulant treatments on guar plants. A-vegetative growth and seed yield, *Sci. J. Flowers Ornam. Plants.* 2017; 4: 143–157.
- [35] Souri MK, Bakhtiarizade M. Biostimulation effects of rosemary essential oil on growth and nutrient uptake of tomato seedlings, *Sci. Hortic.* 2019; 243: 472–476.
- [36] Abbas SM, Akladious SA. Application of carrot root extract induced salinity tolerance in cowpea (*Vigna sinensis* L.) seedlings, *Pak. J. Bot.* 2013; 45: 795–806.